

## **<sup>3</sup>He-rich Solar Particle Mean Free Paths**

J. K. Arballo and B. T. Tsurutani (both at Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109; e-mail: jarballo@jplsp.jpl.nasa.gov, btsurutani@jplsp.jpl.nasa.gov)

G. M. Mason (University of Maryland, College Park, MD 20742)

R. D. Zwickl (NOAA, R/E/SI, 325 Broadway, Boulder, CO 80303)

We have used the largest <sup>3</sup>He-rich flux events of Kahler et al. (1985) to examine potential wave-particle interactions as the particles travel from the Sun to 1 AU. We: 1) search ISEE-3 magnetic field data for in situ waves generated by beam instabilities, 2) examine the particle mfp determined by scattering due to interactions with preexisting "fossil" waves and turbulence, and 3) intercompare the latter with the mfp determined from the particle measurements themselves. Our goal is to attempt to explain why the <sup>3</sup>He-rich events at 1 AU appear to be "scatter-free".

1. 1995 AGU Fall Meeting
2. 001325224
- 3a) B. 'T'. Tsurutani  
Jet Propulsion Laboratory  
MS 169-506  
4800 Oak Grove Drive  
1% adena, CA91109
- b) Tel. 818354-7559
- c) Fax 818354-8895
- d) btsurutani@jplsp.jpl.nasa.gov
4. SPA/SI
5. a)
5. b)
6. Oral
7. 20%, 1993 AGU
8. \$50.00 check
9. C